arXiv:1412.5913v1 [math.AP] 16 Dec 2014

Non-existence of positive weak solutions for some nonlinear $(\mathbf{p},\mathbf{q})$-Laplacian systems

Salah. A. Khafagy*

Mathematics Department, Faculty of Science, Al-Azhar University,
Nasr City (11884), Cairo, Egypt.

E-mail: el_gharieb@hotmail.com

Abstract

In this work we deal with the class of nonlinear $(\mathbf{p},\mathbf{q})$-Laplacian system of the form

$$\left.\begin{array}{rl} -\Delta_p u = \mu\rho_1(x)f(v) & \text{in } \Omega, \\ -\Delta_q v = \nu\rho_2(x)g(u) & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega. \end{array}\right\}$$

where Δ_p with $p > 1$ denotes the p-Laplacian defined by $\Delta_p u \equiv div[|\nabla u|^{p-2}\nabla u]$, μ, ν are positive parameters, $\rho_1(x)$, $\rho_2(x)$ are weight functions, $f, g : [0,\infty) \to \mathbb{R}$ are continuous functions and $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary $\partial\Omega$. Non-existence results of positive weak solutions are established under some certian conditions on f, g when $\mu\nu$ is small.

2000 Mathematics Subject Classification: 35J60,35B40.
Key words: weak solution, p-Laplacian.

1 Introduction:

In this paper we first consider a non-existence result of positive weak solutions for the following nonlinear system

$$\left.\begin{array}{rl} -\Delta_p u = \lambda a_1(x)v^{p-1} - b_1(x)v^{\alpha-1} - c_1(x) & \text{in } \Omega, \\ -\Delta_q v = \lambda a_2(x)u^{q-1} - b_2(x)u^{\beta-1} - c_2(x) & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega, \end{array}\right\} \tag{1.1}$$

where Δ_p with $p > 1$ denotes the weighted p-Laplacian defined by $\Delta_p u \equiv div[|\nabla u|^{p-2}\nabla u]$, $a_i(x)$, $b_i(x)$ and $c_i(x), i = 1, 2$ are weight functions, α and β are positive constants and $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary $\partial\Omega$.

We first show that if $\lambda < \max(\lambda_p, \lambda_q)$, where λ_p, λ_q is the first eigenvalue of $-\Delta_p, -\Delta_q$ respectively, then system (1.1) has no positive weak solutions.

*Current Address: Mathematics Department, Faculty of Science in Zulfi, Majmaah University, Zulfi 11932, P.O. Box 1712, Saudi Arabia.

Next we consider the nonlinear system

$$\left.\begin{array}{rl} -\Delta_p u = \mu\rho_1(x)f(v) & \text{in } \Omega, \\ -\Delta_q v = \nu\rho_2(x)g(u) & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega. \end{array}\right\} \tag{1.2}$$

where μ, ν are positive parameters, $\rho_1(x)$, $\rho_2(x)$ are weight functions and $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary $\partial\Omega$. Let $f, g : [0, \infty) \to \mathbb{R}$ are continuous functions. Also, assume that there exist positive numbers K_i and $M_i, i = 1, 2$ such that

$$f(v) \leq K_1 v^{p-1} - M_1, \quad \text{for all } \ v \geq 0 \tag{1.3}$$

and

$$g(u) \leq K_2 u^{q-1} - M_2, \quad \text{for all } \ u \geq 0. \tag{1.4}$$

We discuss a non-existence result for system (1.2) when $\mu\nu$ is small.

Problems of the form (1.1) and (1.2) arise from many branches of pure mathematics as in the theory of quasiregular and quasiconformal mappings (see [13]) as well as from various problems in mathematical physics notably the flow of non-Newtonian fluids. The p-Laplacian also appears in the study of torsional creep (elastic for $p = 2$, plastic as $p \to \infty$, (see [5]), glacial sliding ($p \in (1; \frac{4}{3}]$, see [10] or flow through porous media ($p = \frac{3}{2}$, see [11]). For existence and non-existence results of positive weak solutions for systems involving the weighted p-Laplacian, see ([2, 3, 6, 7, 8, 9, 12]).

This paper is organized as follows: In section 2, we introduce some technical results and notations, which are established in [4]. In section 3, we prove the non-existence of positive weak solutions for system (1.1) and (1.2).

2 Technical Results

Let us introduce the Sobolev space $W^{1,p}(\Omega)$, $1 < p < \infty$, defined as the completion of $C^\infty(\Omega)$ with respect to the norm (see [4])

$$\|u\|_{W^{1,p}(\Omega)} = \left[\int_\Omega |u|^p + \int_\Omega |\nabla u|^p\right]^{\frac{1}{p}} < \infty. \tag{2.1}$$

Since we are dealing with the Dirichlet problem, we define the space $W_0^{1,p}(\Omega)$ as the closure of $C_0^\infty(\Omega)$ in $W^{1,p}(\Omega)$ with respect to the norm

$$\|u\|_{W_0^{1,p}(\Omega)} = \left[\int_\Omega |\nabla u|^p\right]^{\frac{1}{p}} < \infty, \tag{2.2}$$

which is equivalent to the norm given by (2.1). Both spaces $W^{1,p}(\Omega)$ and $W_0^{1,p}(\Omega)$ are well defined reflexive Banach Spaces.

Now, we introduce some technical results concerning the eigenvalue problem

$$\left.\begin{array}{rl} -\Delta_p u = \lambda a(x)|u|^{p-2}u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{array}\right\} \tag{2.3}$$

We will say $\lambda \in R$ is an eigenvalue of (2.3) if there exists $u \in W_0^{1,p}(\Omega)$, $u \neq 0$, such that

$$\int_\Omega |\nabla u|^{p-2}\nabla u \nabla \phi dx = \lambda \int_\Omega a(x)u^{p-2}u\phi dx, \tag{2.4}$$

holds for $\phi \in W_0^{1,p}(\Omega)$. Then u is called an eigenfunction corresponding to the eigenvalue λ.

Lemma 1 *There exists the first eigenvalue $\lambda_p > 0$ and precisely one corresponding eigenfunction $\phi_p \geq 0$ a.e. in Ω of the eigenvalue problem (2.3). Moreover, it is characterized by*

$$\lambda_p = \frac{\int_\Omega |\nabla \phi_p|^p}{\int_\Omega a(x)|\phi_p|^p} = \inf_{u \in W_0^{1,p}(\Omega)} \frac{\int_\Omega |\nabla u|^p}{\int_\Omega a(x)|u|^p} \leq \frac{\int_\Omega |\nabla u|^p}{\int_\Omega a(x)|u|^p} = \lambda.$$

Definition 1 *A pair of non-negative functions $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ are called a weak solution of (1.2) if they satisfy*

$$\begin{aligned} \int_\Omega |\nabla u|^{p-2}\nabla u \nabla \zeta dx &= \mu \int_\Omega \rho_1(x) f(v)\zeta dx, \\ \int_\Omega |\nabla v|^{q-2}\nabla v \nabla \eta dx &= \nu \int_\Omega \rho_2(x) g(u)\eta dx, \end{aligned}$$

for all test functions $\zeta \in W_0^{1,p}(P, \Omega), \eta \in W_0^{1,q}(\Omega)$.

3 Non-existence Results

In this section we state our main results. Throught this section, we assume q be such that $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 2 *For $\lambda \leq \lambda^*$, system (1.1) has no positive weak solution.*

Proof. Assume that there exist a positive solution $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ of (1.1). Multiplying the first equation of (1.1) by u, we have

$$\begin{aligned} \int_\Omega |\nabla u|^p dx &= \int_\Omega [\lambda a_1(x)v^{p-1} - b_1(x)v^{\alpha-1} - c_1(x)]u dx \\ &< \int_\Omega [\lambda a_1(x)v^{p-1} - c_1(x)]u dx. \end{aligned} \tag{3.1}$$

But, from the characterization of the first eigenvalue, we have

$$\lambda_p \int_\Omega a(x)|u|^p \leq \int_\Omega |\nabla u|^p. \tag{3.2}$$

Combining (1) and (3.2), we have

$$\lambda_p \int_\Omega a_1(x)u^p < \int_\Omega \lambda a_1(x)v^{p-1}u dx - \int_\Omega c_1(x)u dx. \tag{3.3}$$

Similarly, from the second equation of (1.1), we obtain

$$\lambda_q \int_\Omega a_2(x)v^q < \int_\Omega \lambda a_2(x)u^{q-1}v dx - \int_\Omega c_2(x)v dx. \tag{3.4}$$

Adding (3.3) and (3.4), we get

$$\begin{aligned}\lambda_p \int_\Omega a_1(x)u^p + \lambda_q \int_\Omega a_2(x)v^q &< \int_\Omega \lambda a_1(x)v^{p-1}udx + \int_\Omega \lambda a_2(x)u^{q-1}vdx \\ &\quad - \int_\Omega c_1(x)udx - \int_\Omega c_2(x)vdx \\ &< \int_\Omega \lambda a_1(x)v^{p-1}udx + \int_\Omega \lambda a_2(x)u^{q-1}vdx\end{aligned}$$

Applying the Young inequality on the right hand side of the above equation, we have

$$\lambda_p \int_\Omega a_1(x)u^p + \lambda_q \int_\Omega a_2(x)v^q < \int_\Omega \lambda a_1(x)[\frac{u^p}{p} + \frac{v^p}{q}]dx + \int_\Omega \lambda a_2(x)[\frac{v^q}{q} + \frac{u^q}{p}]dx \tag{3.5}$$

Now, we discuss the following two cases:

Case I, if $u \leq v$ for all x, then (3.5) becomes

$$\lambda_p \int_\Omega a_1(x)v^p + \lambda_q \int_\Omega a_2(x)v^q < \int_\Omega \lambda a_1(x)v^p dx + \int_\Omega \lambda a_2(x)v^q dx.$$

Hence,

$$(\lambda_p - \lambda) \int_\Omega a_1(x)v^p + (\lambda_q - \lambda) \int_\Omega a_2(x)v^q < 0$$

which is a contradiction if $\lambda \leq \max(\lambda_p, \lambda_q) = \lambda^*$.

Case II, if $u \geq v$ for all x, then (3.5) becomes

$$\lambda_p \int_\Omega a_1(x)u^p + \lambda_q \int_\Omega a_2(x)u^q < \int_\Omega \lambda a_1(x)u^p dx + \int_\Omega \lambda a_2(x)u^q dx.$$

Hence,

$$(\lambda_p - \lambda) \int_\Omega a_1(x)u^p + (\lambda_q - \lambda) \int_\Omega a_2(x)u^q < 0$$

which is a contradiction if $\lambda \leq \max(\lambda_p, \lambda_q) = \lambda^*$. The proof complete.

Now we consider the main result for system (2.2):

Theorem 3 *Let (1.3) and (1.4) hold. Then system (1.2) has no positive weak solution if* $\mu\nu \leq \frac{\lambda_1^2}{K_1 K_2}$.

Proof. Suppose $u > 0$ and $v > 0$ be such that (u, v) is a solution of (2.2). We prove our theorem by arriving at a contradiction. Multiplying the first equation in (2.2) by a positive eigenfunction say ϕ_p corresponding to λ_p, we obtain

$$-\int_\Omega \Delta_p u \ \phi_p dx = \mu \int_\Omega \rho_1(x)f(v)\phi_p dx,$$

and hence using (2.1) and (1.3), we have

$$\lambda_p \int_\Omega \rho_1(x)u^{p-1}\phi_p dx \leq \mu \int_\Omega \rho_1(x)[K_1 v^{q-1} - M_1]\phi_p dx. \tag{3.6}$$

Similarly using the second equation in (2.2) and (1.4) we obtain

$$\lambda_q \int_\Omega \rho_2(x) v^{q-1} \phi_q dx \leq \nu \int_\Omega \rho_2(x)[K_2 u^{p-1} - M_2]\phi_q dx. \tag{3.7}$$

From (3.7), we have

$$v^{q-1} \leq \frac{\nu}{\lambda_q}[K_2 u^{p-1} - M_2] \tag{3.8}$$

Combining (3.6) and (3.8) we obtain

$$[\lambda_p - \mu\nu\frac{K_1 K_2}{\lambda_q}] \int_\Omega \rho_1(x) u^{p-1}\phi_p \leq -\mu \int_\Omega \rho_1(x)[\frac{\nu K_1 M_2}{\lambda_q} + M_1]\phi_p < 0.$$

Hence system (1.2) has no positive weak solution if $\mu\nu \leq \frac{\lambda_p \lambda_q}{K_1 K_2}$.

Remark 4 *If f, g be such that*

$$f(v) \geq K_1 v^{q-1} + M_1, \quad \text{for all } v \geq 0, \tag{3.9}$$

and

$$g(u) \geq K_2 u^{p-1} + M_2, \quad \text{for all } u \geq 0, \tag{3.10}$$

then we have the following theorem:

Theorem 5 *Let (3.9) and (3.10) hold. Then system (1.2) has no positive weak solution if $\mu\nu \geq \frac{\lambda_p \lambda_q}{K_1 K_2}$.*

Proof. The proof proceeds in the same way as for Theorem 6.

Remark 6 *When $p = q$, $m_1(x) = m_2(x) = m, m = a, b, c$ is constant , and $\alpha = \beta$, we have some results for (1.1) in [1].*

Remark 7 *When $p = q$ and $\rho_1(x) = \rho_2(x) = 1$, we have some results for (1.2) in [1] .*

Acknowledgement. The author would like to express his gratitude to Professor H. M. Serag (Mathematics Department, Faculty of Science, AL- Azhar University) for continuous encouragement during the development of this work.

References

[1] Afrouzi, G. and Valinejad, Z., Nonexistence of result for some p-Laplacian systems, 3(2) (2011) 112-116.

[2] Cirstea, F., Motreanu, D. and Radulescu, V., Weak solutions of quasilinear problems with nonlinear boundary condition, Nonlinear Analysis 43 (2001) 623–636.

[3] P. Drabek and J. Hernandez, Existence and uniqueness of positive solutions for some quasilinear elliptic problem, Nonlinear Anal. 44 (2001) 189–204.

[4] P. Drabek, A. Kufner and F. Nicolosi, Quasilinear elliptic equation with degenerations and singularities, Walter de Gruyter, Bertin, New York, 1997.

[5] Kawohl, B., On a family of torsional creep problems, J. Reine Angew. Math. 410 (1990) 1–22.

[6] Khafagy, S., Existence and uniqueness of weak solution for weighted p-Laplacian Dirichlet problem, Journal of Advanced Research in Dynamical and Control Systems, Vol. 3 (3) (2011), 41– 49.

[7] Khafagy, S., On positive weak solution for a nonlinear system involving weighted p-Laplacian, Journal of Advanced Research in Dynamical and Control Systems, 4 (2012), no. 4, 50-58.

[8] Khafagy, S., Existence and nonexistence of positive weak solutions for a class of weighted (p,q)-Laplacian nonlinear system, Global Journal of Pure and Applied Mathematics, Vol. 9 (4) (2013), 379-387.

[9] Khafagy, S. and Serag, H., Existence of Weak Solutions for $n \times n$ Nonlinear Systems Involving Different p-Laplacian Operators, Electron. J. Diff. Eqns., Vol. 2009, No. 81 (2009) 1-14.

[10] Pélissier, M., and Reynaud, M. Etude d'un modèle mathématique d'écoulement de glacier, C.R. Acad. Sci. Paris Ser. I Math. 279 (1974) 531–534.

[11] Showalter, R. and Walkington, N., Diffusion of fluid in a fissured medium with microstructure, SIAM J. Math. Anal. 22 (1991) 1702–1722.

[12] Serag, H and Khafagy, S., Existence of Weak Solutions for Nonlinear Systems Involving Degenerated p-Laplacian Operators, Sarajevo Journal of Mathematics, Vol. 5, No. 1 (2009) 41-54.

[13] P. Tolksdorf, regularity for a more general class of quasilinear elliptic equations, J. Diff. Eqns., Vol. 51 (1984) 126-150.